

· Drug Category : Antitussives & Expectorants

CODAEWON Tab.

■ COMPOSITION

Each tablet contains Dihydrocodeine bitartrate 5mg, Guaifenesin 50mg, DL- Methylephedrine HCl 17.5mg and Chlorpheniramine maleate 1.5mg.

■ INDICATIONS

Cough, Sputum

■ DOSAGE & ADMINISTRATION

1. Adults : 2 tablets orally, 3 times a day.
2. Children : 8 to 11 years of age - 1 tablet orally, 3 times a day.
11 to 15 years of age - 1 to 2 tablets orally, 3 times a day.

■ PRECAUTIONS

1. The following patients should be advised by a physician or pharmacist prior to taking medicine.
 - 1) Patients who themselves, their parents or brothers are constitutionally predisposed to urticaria, contact dermatitis, allergic rhinitis, migraine, food allergy, etc.
 - 2) Patients with a history of drug allergy (e.g., fever, eruption, joint pain, itching, etc.)
 - 3) Patients with hepatic, renal or thyroid disease, diabetes mellitus, etc.
 - 4) The debilitated.
 - 5) Patients with high fever
 - 6) Patients with cardiac disease, hypertension or the allergy
 - 7) Pregnant or suspected pregnant women or nursing mothers
 - 8) Patients being treated by physician
2. Precautions in administration
 - 1) The advised dosage & administration should be followed.
 - 2) It should not be administered concomitantly with other antitussives & expectorants, cold remedy, antihistamines, sedatives, etc.
 - 3) It should be administered with caution to the children.
3. Precautions during & after administration.
 - 1) When abnormalities (e.g., eruption, rash, nausea, vomiting, anorexia, dizziness, etc.) are observed, discontinue the therapy and take appropriate measures.
 - 2) As drowsiness may occur, the patients should be warned not to engage in activities requiring mental alertness such as operating a motor vehicle or other machinery.

- 3) When symptoms are not improved after the repeated use, discontinue the therapy and should be advised by a physician or pharmacist.
4. Precautions in storage and treatment
 - 1) Keep out of reach of children.
 - 2) Protect from light and store at low humid, cool place.
 - 3) Should not substitute other container.

Caution. This may cause drowsiness. If affected, do not drive or operate machinery.

■ **STORAGE** : Tight container

■ **USE TERM** : 3 years

■ **PACKS** : 30, 100, 500 Tablets

※ Medicine : Keep out of reach of children.

This drug is manufactured in accordance with Korea Good Manufacturing Practice(KGMP) as recommended by WHO.