

Helpovin Inf.

(L-Ornithine-L-Aspartate 5g)

■ DESCRIPTION

1. COMPOSITION

Each ampoule(10mL) contains 5g of L-Ornithine-L-Aspartate (Main material), 1g of D-sorbitol(supplement), Potassium metabisulfite (anti-coloring agent).

2. FORM & COLOR

Colorless, transparent liquid in brown ampoule.

■ INDICATION

Detoxication of symptoms in following disease : Liver cirrhosis, Acute hepatitis, Chronic hepatitis, Hepatic encephalopathy (including pre-coma and coma)

■ DOSAGE & ADMINISTRATION

1. Acute hepatitis : 10-20ml(1-2 ampoules) a day
2. Chronic hepatitis : 20-40ml(2-4 ampoules) a day by drip infusion.
3. Liver cirrhosis : When symptoms are moderate, 10ml(1 ampoule) a time, 10-40ml(1-4 ampoules) a day by drip infusion. When symptoms are severe, up to 80ml(8 ampoules) a day by drip infusion.
4. Liver encephalopathy(including pre-coma, coma) : Up to 80ml (8 ampoules) a day by drip infusion.

Helpovin Inf. may be added to all conventional infusion solutions. However, for reasons of venous tolerance, the dose should not exceed 6 ampoules per 500ml infusion.

■ PRECAUTION

1. Warning

Fragments of glass can drop into a ampoule when ampoule is cut for use, must be careful when you cut ampoule especially use for children and elders.

2. Do not use for patients with following diseases.

Lactic acidosis, Methanol intoxication, Fructose-sorbitol intolerance, Fructose 1.6.-diphosphatase deficiency, sever renal failure.

3. Adverse events

Nervous system : Occasionally, occipital fever sense or nausea may occur.

4. General precautions

Before using high dose, Check Urea value in blood and urine.

5. Precaution for storage

■ **STORAGE** : Hermetic container at room temperature.

■ **USE TERM** : 3 years.

■ **PACKAGE** : 10A/Box

※ **Medicine** : Keep out of reach of children.

This drug is manufactured in accordance with Korea Good Manufacturing Practice (KGMP) as recommended by WHO.