

METHYLERGOMETRINE MALEATE

MERGOT

200mcg/mL Solution for Injection
(IM/IV)

OXYTOCICS

- **FORMULATION** : Each mL contains
Methylergometrine maleate
equivalent to 200mcg
- **INDICATIONS**
 1. Major Indications
For routine management after delivery of the placenta, post-partum atony, hemorrhage and uterine subinvolution.
 2. It can be used after caesarean section and hemorrhage after abortion.
- **DOSAGE AND ADMINISTRATION**

100-200mcg as Methylergometrine maleate at once is IM or IV only in urgency. It may be administered IM as required at intervals of 6 hours, and may be repeated as required at intervals of 2-4 hours in severe hemorrhage.
- **PRECAUTIONS**
 1. Pregnancy
 2. Before delivery of the anterior shoulder
 3. Patients with hypertension and toxemia of pregnancy
 4. Patients with hypersensitivity to ergot alkaloids
 - Should be administered cautiously to the following patients
 - 1) Patients with heart disease or obliterative vascular disease
 - 2) Patients with hepatic or renal disease
 - 3) Patients with sepsis
 - **ADVERSE REACTIONS**
 - 1) Gastrointestinal : Nausea, vomiting, abdominal pain, diarrhea, etc. may occur.
 - 2) Cardiovascular : Hypertension, tachycardia, bradycardia, etc. may occur.
 - 3) Psychoneurotic : Headache, drowsiness, dizziness, dry mouth, tinnitus, excitation, etc. may occur.
 - 4) Miscellaneous : Chest pressure sensation, etc. may occur.
- **PRECAUTIONS IN APPLICATION**
 - 1) This drug should be injected slowly with careful monitoring of blood pressure when it is used intravenously concomitantly with anesthetics, hypertensors, etc.
 - 2) In case of intramuscular injection, be careful of the following to avoid affecting the nerve, tissue, etc...
 - a) Be careful to avoid injecting on the nerve track area. In case of scaly pain, pull the needle out and insert the needle on another site.
 - b) In case of repeated injection, substitute the injection sites. (Avoid repeated injection to the same site).
- **CAUTION**

Foods, Drugs, Devices and Cosmetics Act prohibits dispensing without prescription.
- **STORAGE**

Store at temperatures between 15°C to 25°C. Protect from light.
- **AVAILABILITY** : 50 Ampoules/Box, 10 Ampoules/Box