



Korean Ginseng + Royal Jelly

Oramin-F

Soft Cap.

■ COMPOSITION : Each soft capsule contains

Korean ginseng	50mg
Royal Jelly	10.0mg
Retinol palmitate (Vitamin A)	2500 I.U.
Ergocalciferol (Vitamin D ₂)	400 I.U.
dl- α -Tocopherol acetate (Vitamin E)	45mg
Thiamine nitrate (Vitamin B ₁)	2mg
Riboflavin (Vitamin B ₂)	2mg
Pyridoxine hydrochloride (Vitamin B ₆)	2mg
Ascorbic acid (Vitamin C)	100mg
Nicotinamide (Vitamin B ₃)	20mg
Calcium pantothenate (Vitamin B ₅)	30mg
Cyanocobalamin (Vitamin B ₁₂)	6.0 μ g
Ferrous fumarate	27.38mg
Copper sulfate	7.86mg
Magnesium oxide	66.34mg
Zinc oxide	5.0mg
Anhydrous dibasic calcium phosphate	307.5mg
Manganese sulfate	3.0mg
Potassium sulfate	18.0mg

■ INDICATIONS

1. Supplementation of vitamin A, B₁, B₂, B₆, C, D, E in following cases :
Fatigue, pregnancy, the lactation, weak constitution, the period of growth, senescence.
2. Alleviation of xerophthalmos, night blindness
3. Ateliosis of bone and teeth, prevention of rickets

■ DOSAGE & ADMINISTRATION

The usual dosage for Children over age 12 and adults is 1 soft capsule daily.

■ CAUTIONS

1. Warnings.

- 1) In pregnancy, administration of Vit. A(Retinol) more than 5,000 I.U. may cause congenital abnormality.
(Except patients with Vitamin A deficiency).
- 2) In case of administration to under 6-year children may cause toxicity.

2. Do not administer to the following patients.

- 1) Patients with hypercalcemia, hyperkalemia sarcoidosis, nephrolithiasis, severe renal failure, Wilson disease, hematochromatosis.
- 2) Infants under 1 year of age.
- 3) Patients with hypersensitivity with this product.

3. Should be administered cautiously to the following patients.

(Consult a doctor or a pharmacist)

- 1) Patients undergoing therapy.
- 2) Patients with hyperoxaluria, renal disorder, hypoproteinos, cardiovascular dysfunction, circulatory failure.
- 3) Patients with gastrointestinal disease(ex)Peptic ulcer, chronic ulcerative colitis, regional colitis)
- 4) Patients with cardiac glycoside administration.

4. Adverse reactions.

- 1) Cease the administration with following symptom and Consult a doctor or a pharmacist.
 - Pruritus, eruption, flare, erythema, xeroderma, nausea, vomit, diarrhea, constipation, epigastric distress, anorexia, abdominal distention, epigastralgia, photosensitivity, convulsant, fever, coma, asystole, CNS dysfunction, hyperkalemia, pulmonary edema, urticaria.
- 2) Oramin-F may cause acceleration of menstruation, menorrhagia, extension of anovulatory menstrual cycle.
- 3) Oramin-F may cause thrombosis to Patients with oral contraceptive administration or thrombotic predisposition.
- 4) Mass administration may cause nausea, vomit, hypernatremia, congestive heart failure, edema.
- 5) Long-term administration may cause hypercalcemia, nephrolithiasis.

5. General precautions

- 1) If you think that the product is having too little effect in 1 month, you should discuss the matter with a doctor or a pharmacist.
- 2) Do not take more than 2 capsules of Oramin-F. (Do not take more than 5,000IU of retinol a day as supplement)

6. Contraindications.

- 1) Combination therapy with Aldosterone antagonist and triamterene may cause hypercalcemia.
- 2) Patients taking oral tetracyclines, antacid, phosphates, Levo-dopa.
- 3) Do not administer before and after drinking teas contain tannins.

7. Cautions in keeping and handling

- 1) Keep out of reach of children.
- 2) Store at cool places, protected from a direct ray of light.
- 3) Do not preserve in another containers in order to maintain quality of the product and avoid misuse.

■ **STORAGE** : Preserve in Air-tight container, protected from light.

■ **USE TERM** : 3 years from MNF. Date

■ **PACKS** : 30, 60 Caps/Box

Manufactured by

Daewon Daewon Pharm.Co.,Ltd. **KGMP**

903-1, Sangshin-ri, Hyangnam - myun, Hwaseong, Kyunggi - do, Korea