

# Daewon PRODUCT LIST





The most important value  
for us is belief and trust.

#### History

- 1958. 01. Founded
- 1988. 12. Completed Hyangnam KGMP factory construction
- 1989. 09. Established the Central Research Laboratory
- 1994. 10. Listed on the Korea Securities Dealer Automated Quotation (KOSDAQ)
- 1999. 12. Listed on the Korea Stock Exchange (KOSPI)
- 2007. 04. Acquired approval of Pelubi Tablet as the 12th new drug developed in Korea
- 2008. 01. 50th anniversary of foundation
- 2009. 02. Awarded at the 10th Korea New Drug Development Award (Technology Prize; Pelubi Tab.)
- 2012. 06. Certified as an Innovative Pharmaceutical Enterprise (Ministry of Health and Welfare)
- 2015. 07. Designated as an Advanced Technology Center (Ministry of Commerce, Industry and Energy)
- 2015. 12. Honored Grand Prize of Good Company Award (Citizens' Coalition for Economic Justice)
- 2016. 02. Awarded at the 17th Korea New Drug Development Award (Technology Prize; Pelubi CR Tab.)
- 2018. 06. ISO 37001 Approved
- 2020. 01. Establishment of New Factory in Jincheon
- 2021. 05. Acquired Keukdong H Pharm (transformed to Daewon Healthcare)

#### Factory



- Jincheon Factory
- Plottage : 79,922 m<sup>2</sup>
- Capacity : 580 million of liquid dosage unit (Biggest capacity in Korea)



- Hyangnam Factory
- Plottage : 15,269 m<sup>2</sup>
- Capacity : 2 billion of solid dosage unit  
25 million of liquid dosage unit  
100 million of injectable dosage unit



## Strength of Liquid Stick Sachet Specialized in CMO Business

We, Daewon Pharm., are specialized  
in manufacture of Liquid Stick Sachet.


#### Portable, Convenient and Fast!

- Increased adherence of patients
- More than 81% of repurchase rate
- Possible to formulate diverse pharmaceuticals & food supplements



#### Expand to Overseas Business

Daewon Pharm is strategically positioning itself for robust expansion, with active operations including a subsidiary in Vietnam. Our approach involves licensing out advanced medications including the new drug such as Pelubi & Pelubi CR Tablets (NSAIDs), Megex-I Suspension (anti-cancer supplement), and Privituss Suspension (antitussive), with a particular focus on pharmerging markets. Our commitment extends to exporting cutting-edge technologies and distinct pharmaceutical offerings, all rooted in our proprietary platform technology. Additionally, we are actively exporting health functional supplements such as Oramin series (Korean Ginseng Vitamin) and probiotics.



#### Our vision heads to total health care company by acquiring Daewon Healthcare.


We acquired Daewon Healthcare as an affiliate specialized in manufacturing health functional supplement. Daewon Healthcare offers finished supplements such as multi-vitamin, probiotics and contract manufacture of customer brand.






Product	Active ingredient	Indication	Dosage & Administration
---------	-------------------	------------	-------------------------


Pelubi Tab. Analgesic

	Each tablet contains <b>Pelubiprofen 30mg</b>	Reduce symptoms of Osteoarthritis, Rheumatoid arthritis, Back pain (Lumbago) and Fever	1 Tab.(30mg), 3 times daily
---	--	---	-----------------------------


Pelubi CR Tab. Analgesic

	Each tablet contains <b>Pelubiprofen 45mg</b>	Reduce symptoms of Osteoarthritis, Rheumatoid arthritis, Back pain (Lumbago) and Acute pain	1 Tab. (45mg), twice daily.
---	--	--	-----------------------------

Curefen Syr. Analgesic

	In 1mL <b>Dexibuprofen 6mg, 12mg</b>	Antipyretic analgesic	The doses are different according to age.
---	---	-----------------------	--


Senafen Tab. Analgesic

	Each tablet contains <b>Aceclofenac 100mg</b>	Rheumatoid arthritis, Ankylosing spondylitis, Osteoarthritis	1 Tab., twice daily
---	--	--	---------------------


Wontran ER Tab. Analgesic

	Each tablet contains <b>Acetaminophen 650mg</b> <b>Tramadol HCl 75mg</b>	Moderate to severe chronic pain	Take 1 Tab. at first dose, administrate 2 Tabs. once, twice daily. (4 Tabs a day) Administration interval should be at least 12 hours. The maximum daily dose of Wontran ER tab is 4 Tabs.
---	--	---------------------------------	---

Nerexone SR Tab. Analgesic


	Each tablet contains <b>Eperisone Hydrochloride 75mg</b>	Painful muscle spasms associated with musculoskeletal disorders: cervical brachial syndrome, shoulder pain, back pain	Adults: 2 times a day, 1 tablet once (75 mg as eperisone hydrochloride) orally administered after meals. Increase or decrease appropriately according to age and symptoms. Take it whole without crushing, dividing, or chewing.
---	---	---	--

Codaewon Tab. Respiratory

	Each tablet contains <b>Dihydrocodeine bitartrate 5mg</b> <b>DL-Methylephedrine HCl 17.5mg</b> <b>Chlorpheniramine maleate 1.5mg</b> <b>Guaifenesin 50mg</b>	Cough, Sputum	Adults : 2 Tabs., 3 times daily Children : 11~15years : 1~2 Tabs. / 8~11years : 1 Tab. three times daily
---	--	---------------	---

Product	Active ingredient	Indication	Dosage & Administration
---------	-------------------	------------	-------------------------


Nezelex Nasal Spray Respiratory

	In 1 dose <b>Mometasone Furoate 50μg</b>	Allergic rhinitis	If patients push the pump one time, Mometasone Furoate 100mg release(Mometasone Furoate monohydrate 0.05mg). Once daily, spraying twice in nasal.
---	---	-------------------	---


Rizine Tab. Respiratory

	Each tablet contains <b>Cetirizine HCl 10mg</b>	Rhinitis by seasonal allergy, Conjunctivitis, Chronic rhinitis, Itching	Over 6 year and adults : 1 Tab., daily, administer in the evening.
---	--	---	---


Xyzine Tab. Respiratory

	Each tablet contains <b>Levocetirizine HCl 5mg</b>	Seasonal and perennial allergic rhinitis, the uncomplicated skin manifestations of chronic idiopathic urticaria	Adults and children over 6 years of age : 1 Tab.(Levocetirizine 5mg), once daily
---	---	--	--


Codaewon Forte Syr. Respiratory

	In 100ml syrup <b>Ammonium Chloride 1000mg</b> <b>Chlorpheniramine Maleate 15mg</b> <b>Dihydrocodeine Tartrate 50mg</b> <b>DI-methylephedrine HCl 131mg</b>	Antipyretic analgesic	The doses are different according to age.
---	---	-----------------------	--


Codaewon S Syr. Respiratory








	In 1mL <b>Dihydrocodeine Tartrate Hydrochloride 10mg</b> <b>DL-Methylephedrine Hydrochloride 26.2mg</b> <b>Chlorpheniramine Maleate 3mg</b> <b>Ammonium Chloride 200mg</b> <b>Pelargonium Sidoides 11%</b> <b>Ethanol Extract (1→8~10) 1235.52mg</b>	Improvement of symptoms and signs of acute bronchitis	Take 1 packet (20 mL) once for adults, 3 times a day.
---	--	--	--








Beposta B Tab. Respiratory

	Each tablet contains <b>Bepotastine besilate 10mg</b>	Treatment of itching due to allergic conjunctivies	1 Tab., twice daily
---	--	---	---------------------









Elstein Tab. / Cap. / Syr. Respiratory







	Each tablet, capsule and syrup contains <b>Erdosteine 300mg, 300mg and 35mg/mL</b>	Relief of cough acute & chronic respiratory disease	Tablet and Capsule : 1 Cap, 2~3 times daily Syrup : 175~350mg, once or twice daily
---	--	--	---








Product	Active ingredient	Indication	Dosage & Administration
Oseltawon Cap. <span>Respiratory</span>			
	Each capsule contains <b>Oseltamivir phosphate 30mg, 45mg, 75mg</b>	Treatment of Influenza	In adults and adolescents 13 years and older is 75 mg twice daily for 5 days.
Trigel Susp. <span>Gastrointestinal</span>			
	Each pouch(10mL) contains <b>Oxethazaine 20mg</b> <b>Dried Aluminum hydroxide gel 582mg</b> <b>Magnesium hydroxide 196mg</b>	Acute & Chronic pain, Gastritis, Duodenal ulcer, Esophagitis, Stomach pain by dyspepsia in radiotherapy	1/2~1 Pouch, 4 times daily
Eswonamp Tab. <span>Gastrointestinal</span>			
	Each tablet contains <b>Esomeprazole magnesium 20, 40mg</b>	GERD, Zolonger-Elison syndrome, Duodenal ulcer	GERD : 20~40mg daily 20mg or 40mg Duodenal Ulcer : 20mg daily
Escorten Tab. <span>Gastrointestinal</span>			
	Each tablet contains <b>Esomeprazole Magnesium Trihydrate 11.15mg</b>	Improvement of gastric mucosal lesions in acute gastritis and chronic gastritis	Adults: 1 tablet at a time, orally administered once a day. Children under 12 years of age: There is no case of administering this drug to children under 12 years of age.
Fotagel Susp. <span>Gastrointestinal</span>			
	Each pouch(20mL) contains <b>Diocahedral smectite 3g</b>	Pain symptoms after esophageal, gastroduodenal and enteric disease. Acute or chronic diarrhea.	Acute : 20mL, 3~4 times daily Children : under 6 months of age : 10mL, 2 times daily 6 months~5 years : 10mL, 3 times daily over 5 years : 20mL, 2~3 times daily
Freegel Susp. <span>Gastrointestinal</span>			
	Each pouch(15mL) contains <b>Almagate 1.5g</b>	Gastroduodenal ulcer, Gastritis, Hyperchylia (heartburn, nausea, vomiting, gastric pain, belching up a bit of sour vomit), Reflux esophagitis.	1 Pouch(15mL) 3 times daily, in 1~2 hours after meals. In case of severe gastroduodenal ulcer, 1 Pouch(15mL) 4 times daily in 1~2 hours after meals or before sleeping.
Pantyrase Tab. <span>Gastrointestinal</span>			
	Each tablet contains <b>Pancreatin 175mg</b> <b>Ox bile Extract 25mg</b> <b>Dimethicone 25mg</b> <b>Hemicellulase 50mg</b>	Dyspepsia, Anorexia, Abdominal fullness	Adults : 1~2 Tabs., 3 times daily








Product	Active ingredient	Indication	Dosage & Administration
Pariben Tab. <span>Gastrointestinal</span>			
	Each tablet contains <b>Rabeprazole sodium 10mg, 20mg</b>	Gastric ulcer, Duodenal ulcer, Oesophagitis erosiva	Gastric ulcer, Duodenal ulcer : 10mg once daily, Oesophagitis erosive : 20mg once daily
Clasin dried syr. 125mg/5mL, 250mg/5mL <span>Antimicrobial</span>			
	125mg/5mL : Each 1mL contains <b>Clarithromycin 25mg(activity)</b>  250mg/5mL : Each 1mL contains <b>Clarithromycin 50mg(activity)</b>	Upper & lower respiratory infection, skin infection, Mycobacterium infection	7.5mg/kg(as clarithromycin activity), Every 12 hours
Clasin Tab. <span>Antimicrobial</span>			
	Each tablet contains <b>Clarithromycin 250, 500mg</b>	Lower & Upper respiratory tract infections: bronchitis, pneumonia, pharyngitis, sinusitis, etc.	Adults: As clarithromycin, 250mg (potency) is usually administered twice a day, and in case of severe infection, 500mg (potency) is administered twice a day.
Wonfuroxime Inj. <span>Antimicrobial</span>			
	Each vial contains <b>Cefuroxime sodium 250mg, 750mg, 1.5g</b>	Bronchitis, Pneumonia, Pyelonephritis, Meningitis, Gonorrhea	Adults : 750mg 3 times daily, IV or IM Children : 30~100mg / kg divided into 3~4 times daily IV or IM
Wontaxime Inj. <span>Antimicrobial</span>			
	Each vial contains <b>Cefotaxime sodium 250mg or 500mg or 1g</b>	Bronchitis, Pneumonia, Meningitis, Gonorrhea, Septicemia, Osteomyelitis, Peritonitis	Acute : 2~12g divided into several times IV or IM according to infections Children: 50~100mg / kg divided every 6~12 hours daily IV
Wontiaxone Inj. <span>Antimicrobial</span>			
	Each vial contains <b>Ceftriaxone sodium 1g, 0.5g</b>	Various infections	Adult : 1~2g IV and IM once daily. In severe infection up to 4g may be given Pediatric : 20~50mg/kg IV and IM once daily. In severe infection up to 80mg/kg may be given
A-phine Tab. <span>Cardiovascular</span>			
	Each tablet contains <b>Amlodipine maleate 6.42mg (5mg as Amlodipine)</b>	Hypertension, stable-angina pectoris	The doses are different according to patients. 1 Tab., once daily (maximum dose/day : 10mg)










Product	Active ingredient	Indication	Dosage & Administration
Clow Tab. <span>Cardiovascular</span>	 Each tablet contains Rosuvastatin 5mg, 10mg, 20mg	Protopathy hypercholesterolemia, Combined hyperlipidemia	5~20mg, once daily
Clow Z Tab. <span>Cardiovascular</span>	 Each tablet contains Rosuvastatin 5mg, 10mg, 20mg Ezetimibe 10mg	Protopathy hypercholesterolemia, Combined hyperlipidemia	1 Tab., once daily
Litozet Tab. <span>Cardiovascular</span>	 Each tablet contains Ezetimibe 10mg Atorvastatin 10mg, 20mg, 40mg	Primary hypercholesterolemia	This drug is administered once daily, regardless of meal
Diortan Susp. <span>Cardiovascular</span>	 Each tablet contains Valsartan 80mg, 160mg	Hypertension	1 Tab., once daily
Daewon Clopidogrel Tab. <span>Cardiovascular</span>	 Each tablet contains Clopidogrel Bisulfate 97.875mg (75mg as Clopidogrel)	Recent MI, Recent Stroke or Established Peripheral Arterial Disease. Acute Coronary Syndrome. Decrease the risk of atherothrombosis and thromboembolism.	1 Tab., once daily
Itorin Tab. <span>Cardiovascular</span>	 Each tablet contains Simvastatin 10mg, 20mg Ezetimibe 10mg	Protopathy hypercholesterolemia, Combined hyperlipidemia	The doses are different according to patients
Lipiwon Tab. <span>Cardiovascular</span>	 Each tablet contains Atorvastatin Ca 10.36 / 20.72mg (10, 20mg as Atorvastatin)	Patients with primary hypercholesterolemia and prevention of Cardiovascular Disease	1 Tab.(10mg), once daily. If needed, the maximum dosage is 80mg per day.
Wonsartan Tab. <span>Cardiovascular</span>	 Each tablet contains Losartan potassium 50mg	Hypertension	1 Tab.(50mg), once daily. If needed, the maximum dose is 2 Tabs.(100mg).








Product	Active ingredient	Indication	Dosage & Administration
Wonvelol Tab. <span>Cardiovascular</span>	 Each tablet contains Carvedilol 12.5mg / 25mg	Essential hypertension, Chronic stable angina pectoris	Essential hypertension : 25mg once daily (2 weeks) Chronic stable angina pectoris : 25mg 2times daily (2 weeks)
Twincombi Tab. <span>Cardiovascular</span>	 Each tablet contains Telmisartan 40mg + Amlodipine 5mg Telmisartan 40mg + Amlodipine 10mg Telmisartan 80mg + Amlodipine 5mg	Hypertension	1 Tab., once daily
Tri-in-one Tab. <span>Cardiovascular</span>	 Each tablet contains Telmisartan 40mg + Amlodipine 5mg + Rosuvastatin 5mg Telmisartan 40mg + Amlodipine 5mg + Rosuvastatin 10mg Telmisartan 80mg + Amlodipine 5mg + Rosuvastatin 5mg Telmisartan 80mg + Amlodipine 5mg + Rosuvastatin 10mg	Hypertension & Hyperlipidemia	1 Tab., once daily
TG Fenone Tab. <span>Cardiovascular</span>	 Each tablet contains Choline Fenofibrate 178.8 mg (as fenofibric acid 135 mg)	Treatment of Severe Hypertriglyceridemia and Treatment of Primary Hypercholesterolemia or Mixed Dyslipidemia	1 Tab., once daily
Spiracton Tab. <span>Cardiovascular</span>	 Each tablet contains Spironolactone 25mg, 50mg	Heart failure, Hypertension and hypokalemia	1~2 Tabs., daily
Uptava Cap. <span>Cardiovascular</span>	 Each tablet contains Pitavastatin calcium 2mg, Fenofibrate Granules 241.51mg	Treatment of complex dyslipidemia with high triglyceride levels and low HDL-cholesterol levels, although LDL-cholesterol levels are adequately controlled during pitavastatin 2mg monotherapy in adult patients at high risk of coronary heart disease (CHD)	Take 1 capsule once a day immediately after meals. Serum lipid levels (total cholesterol (TC), LDL-C, and triglycerides (TG)) should be monitored in response to treatment. patients receiving pitavastatin and fenofibrate concomitantly are not switched to this drug since there is no bioequivalence data.

Product	Active ingredient	Indication	Dosage & Administration
Mericle Tab. <span>Antidiabetic</span>	 Each tablet contains <b>Glimepiride 2mg</b>	Insulin-independent diabetes(Type II)	It should be adjusted according to patients. 1 Tab., once daily. It should be administrated before the first meal of a day.
Altiro Tab. <span>Antidiabetic</span>	 Each tablet contains <b>R-Thioctic acid(R-α-Lipoic Acid) 480mg</b>	Diabetic Peripheral Neuropathy	1 Tab., daily
Janulitin Tab. <span>Antidiabetic</span>	 Each tablet contains <b>Sitagliptin 25, 50, 100mg</b>	This drug is administered as an adjunct to diet and exercise therapy to improve blood sugar control in patients with type 2 diabetes.	Take 100 mg once a day in monotherapy or combination therapy, and the maximum daily dose is 100 mg. This medicine can be taken regardless of meals.
Janulitin Combi Tab. <span>Antidiabetic</span>	 Each tablet contains <b>Sitagliptin 50mg Metformin hydrochloride 500, 850, 1000mg</b>	This drug is administered as an adjunct to diet and exercise therapy to improve blood sugar control in patients with type 2 diabetes. If metformin or sitagliptin monotherapy cannot adequately control blood sugar or if sitagliptin and metformin combination therapy is substituted	Take twice a day with food, and the dose is gradually increased to reduce gastrointestinal side effects caused by metformin.
Dapasita Tab. 10/100mg <span>Antidiabetic</span>	 Each tablet contains <b>Dapagliflozin 10mg Sitagliptin 100mg</b>	This drug is administered as an adjunct to diet and exercise to improve blood sugar control in type 2 diabetic patients for whom sitagliptin and dapagliflozin are appropriately administered.	Take 1 tablet daily regardless of meal. Take it whole without crushing, dividing, or chewing.
Dapasita-M SR Tab. <span>Antidiabetic</span>	 Each tablet contains <b>Dapagliflozin 5mg, 10mg Sitagliptin 50mg, 100mg Metformin hydrochloride 500mg, 750mg, 1000mg</b>	This drug is administered as an adjunct to diet and exercise therapy to improve glycemic control in type 2 diabetic patients for whom a combination of dapagliflozin, sitagliptin and metformin is appropriate.	Patients taking 10/100/1000 mg sustained-release tablets of this drug should take 1 tablet once a day. Patients taking 5/50/500 mg, 5/50/750 mg and 5/50/1000 mg extended-release tablets of this drug should take 2 tablets once a day.
Dapawon Tab. <span>Antidiabetic</span>	 Each tablet contains <b>Dapagliflozin 5mg, 10mg</b>	Type 2 Diabetes: This drug is administered as an adjunct to diet and exercise to improve blood sugar control in patients with type 2 diabetes.	This medicine may be taken orally once daily at any time of the day, regardless of the meal. Take it whole without crushing, dividing, or chewing.

Product	Active ingredient	Indication	Dosage & Administration
Dapawon M Tab. <span>Antidiabetic</span>	 Each tablet contains <b>Dapagliflozin 5mg, 10mg Metformin hydrochloride 500mg, 1000mg</b>	Type 2 Diabetes: This drug is administered as an adjunct to diet and exercise to improve blood sugar control in patients with type 2 diabetes.	This drug is usually administered once daily with the evening meal. It should be individualized without exceeding the maximum recommended dose of dapagliflozin 10mg and extended release metformin 2000mg.
Licure Susp. <span>Hepatic</span>	 Each pouch(20g) contains <b>L-leucine 0.952g L-isoleucine 1.904g L-valine 1.144g</b>	Hypoalbuminemia	1 Pouch(20g), 3 times daily
Metran Cap. <span>Psychoneurotic</span>	 Each capsule / ampoule(1mL) contains <b>Mecobalamin 500μg</b>	Peripheral neuronc disorder. Megaloblastic anemia caused by deficiency of Vitamin B12.	Cap. : Adults : 1500μg daily divided into 3 times Inj. : Adults : 1 Amp. / day 3 times a week IM, SC
Alfocholine Soft cap. / Inj. / Syr. <span>Psychoneurotic</span>	 Each Soft cap. / ampoule / pouch contains <b>Choline Alfoscerate 400mg / 1g / 400mg</b>	Improvement of cognitive dysfunction in patients with dementia of neurodegenerative and vascular origin	Tablet and Syrup : 400mg, 2~3 times daily Injection : 1g, once daily
Terasin Tab. <span>Endocrine</span>	 Each tablet contains <b>Terazosin HCl 2.314mg(as Terazosin 2mg)</b>	Hypertension(mild-moderate). Urinary symptoms of benign prostatic hyperplasia	Hypertension: Initial dose 1mg, maintenance dose 2~10mg. BPH: Initial dose 1mg, maintenance dose 5~10mg.
Agnafil Tab. <span>Endocrine</span>	 Each tablet contains <b>Sildenafil 100mg</b>	Impotence	50mg once as needed, approximately 1 hour before sexual activity
Solipro Tab. <span>Endocrine</span>	 Each tablet contains <b>Solifenacin succinate 5mg, 10mg</b>	Overactive bladder with symptoms of urge urinary incontinence, urgency, and frequency	5mg~10mg, once daily




Product	Active ingredient	Indication	Dosage & Administration
Oramin-G Soft Cap. Mutivitamin			
	Each soft capsule contains Korean Ginseng 40mg Aloe extr. 5mg Vitamins Minerals	Supplementation of vitamins and minerals	Adults : 1 Cap., daily
Oramin-F Soft Cap. Mutivitamin			
	Each soft capsule contains Korean Ginseng 40mg Aloe extr. 5mg Vitamins Minerals	Supplementation of vitamins and minerals	Adults : 1 Cap., daily
Oramin-E Soft Cap. Mutivitamin			
	Each soft capsule contains Korean Ginseng 100mg Garlic ext. 10mg Vitamins Minerals	Supplementation of vitamins and minerals	Adults : 1 Cap., daily
Renamezin Cap. Kidney Disease			
	In one time dose (7 capsules) Spherical adsorptive carbon 2g	Chronic renal failure	7 Caps., 3 times daily (21 Caps./ day)
Lowpota Suspension Kidney Disease			
	Each 1 pouch (20 mL) contains Calcium Polystyrene Surfonate 5g	Hyperkalemia	Calcium polystyrene sulfonate, 15 to 30 g per day divided into 2 to 3 doses and administered orally.
Semerwon Tab. Kidney Disease			
	Each tablet contains Sevelamer carbonate 800mg	Serum phosphorus control in patients with chronic kidney disease receiving dialysis	This medicine should be taken three times a day with meals.
Tolvis Tab. Kidney Disease			
	Each tablet contains Tolvaptan 15mg, 30mg	Clinically significant hypervolemic or euvolemic hyponatremia in patients with heart failure, syndrome of abnormal antidiuretic hormone secretion (SIADH), etc	The recommended initial dose for adults is 15 mg once daily. To reach the target serum sodium concentration, increase the dose to 30 mg once a day at an interval of at least 24 hours, and increase to a maximum of 60 mg once a day. To minimize the risk of liver damage, do not administer for more than 30 days.


Product	Active ingredient	Indication	Dosage & Administration
Risewon Plus Tab. <b>Osteoporosis</b>			
	Each tablet contains <b>Risedronate Sodium 2.5 Hydrate 40.162mg + Cholecalciferol(Vitamin D) Concentrated Powder 56mg</b>	Treatment and prevention of osteoporosis in postmenopausal women / Treatment of osteoporosis in men	1 Tab., weekly
Forend Tab. <b>Osteoporosis</b>			
	Each tablet contains <b>Alendronate sodium 91.37mg</b>	Osteoporosis	1 Tab., once weekly
Forend Plus D Tab. <b>Osteoporosis</b>			
	Each tablet contains <b>Alendronate sodium 91.37mg Cholecalciferol concentrated powder 53.33mg(5600IU as Vitamin D3)</b>	Treatment of osteoporosis in postmenopausal woman. Treatment to increase bone mass in men with osteoporosis.	1 Tab., once weekly
Curasten Sol. <b>Others</b>			
	Each 10mL contains <b>Citrulline maleate 1g</b>	Adynamia	10mL, 2~3 times daily
Freefol MCT Inj. 20mL, 50mL <b>Others</b>			
	In 1mL <b>Propofol 10mg (1%) Propofol 20mg (2%)</b>	Anesthesia	General Anesthesia - Adults administration 4mL, every 10sec. for induction.
Megex-I Susp. <b>Others</b>			
	In 1mL <b>Megestrol acetate 40mg</b>	Anorexia, Cachexia of cancer or AIDS patients or predominant weight loss of unknown cause.	For initial dose, 20mL daily for adults
Megex -I ES Susp. <b>Others</b>			
	In 1mL <b>Megestrol acetate 125mg</b>	Anorexia, Cachexia of cancer or AIDS patients/ Significant weight loss caused by unknown reason.	Take Megestrol Acetate 625 mg(5mL)/ day. Shake well before administration.

Product	Active ingredient	Indication	Dosage & Administration
---------	-------------------	------------	-------------------------


Coldwon Cold Q Syr. **OTC**

	Each pouch contains Acetaminophen 325 mg Chlorpheniramine Maleate 2.5 mg Ephedrine HCl 18 mg Guaifenesin 42 mg	Temporarily relieves common cold/flu symptoms	Please see the insert
---	--	---	-----------------------


Coldwon Cough Q Syr. **OTC**

	Each pouch contains Acetaminophen 325mg Dextromethorphan HBr 16mg dl-methylephedrine HCl 21mg Guaifenesin 83mg Caffeine Anhydrous 25mg	Temporarily relieves common cold/flu symptoms	Please see the insert
--	---	---	-----------------------

Coldwon Nose Q Syr. **OTC**


	Each pouch contains Acetaminophen 325mg Pseudoephedrine HCl 30mg Tripolidine Hydrochloride Hydrate 1.33mg Caffeine Anhydrous 25mg	Temporarily relieves common cold/flu symptoms	Please see the insert
---	---	---	-----------------------

Coldwon Kids Cold Syr. **OTC**


	Each pouch contains Acetaminophen 75mg Chlorpheniramine Maleate 0.9375mg Dextromethorphan HBr 6mg dl-methylephedrine HCl 9.375mg Guaifenesin 31.25mg	Temporarily relieves common cold/flu symptoms	Please see the insert
---	---	---	-----------------------

Product	Active ingredient	Indication	Dosage & Administration
---------	-------------------	------------	-------------------------


Coldwon Kids Fen Syr. **OTC**

	Each pouch contains Acetaminophen 160mg	Temporarily relieves common cold/flu symptoms	Please see the insert.
---	--	---	------------------------


Coldwon Kids Ibufen Syr. **OTC**

	Each pouch contains Ibuprofen 100mg	Temporarily relieves common cold/flu symptoms	Please see the insert.
--	--	---	------------------------

Coldwon Kids Cough Syr. **OTC**

	Each pouch contains Guaifenesin 10.5mg Pentoxifyverine citrate 2.5mg DL-Methylephedrine HCl 3.125mg	Temporarily relieves common cold/flu symptoms	Please see the insert.
---	--	---	------------------------

Coldwon Kids Nose Syr. **OTC**

	Each pouch contains Chlorpheniramine Maleate 1.0 mg Pseudoephedrine HCl 15mg	Temporarily relieves common cold/flu symptoms	Please see the insert.
---	--	---	------------------------



Advanced healthcare company, for better life.

**Daewon** Pharm.Co.,Ltd.

E-mail [export@daewonpharm.com](mailto:export@daewonpharm.com) Tel 82-2-2204-6971~9 Fax 82-2-499-9110

Head Office 386 Cheonhodaero, Sungdong-gu, Seoul, Korea

Hyangnam Factory 24 Jeyakgongdan 1-gil, Hyangnam-eup, Hwaseong-si, Gyeonggi-do, Korea

Jincheon Factory 44 Yongso 2-gil, Gwanghyewon-myeon, Jincheon-gun, Chungcheongbuk-do, Korea

DW\_230921\_V.0000